



### General

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_ Height \_\_\_\_\_ Weight \_\_\_\_\_ Phone # \_\_\_\_\_

Order Start Date \_\_\_\_\_ Order Expiration Date \_\_\_\_\_

Allergies: \_\_\_\_\_

### Guidelines for Ordering:

1. Send **FACE SHEET** and **H&P** or most recent chart note.
2. Send pertinent **labs**, such as ferritin level, (must be within last 30 days).

### DIAGNOSIS CODE: \*\*IF using 'Other' must include ICD10 code\*\*

- |   |   |
|---|---|
| <input type="checkbox"/> D50.9 Iron deficiency anemia           | <input type="checkbox"/> D50.0 Iron deficiency anemia due to chronic blood loss |
| <input type="checkbox"/> D63.1 Anemia in chronic kidney disease | <input type="checkbox"/> Other: _____   |

### Nursing Orders

**IV ACCESS:**     Place peripheral IV     Access Implanted Port

**\*If nurse to access implanted port please complete Port Access Order Set as well\***

Vital Signs: baseline and every 30 minutes until discharge. Hold patient for 30 minutes post-infusion to observe/monitor for adverse reaction

### Pre-Medications

- |   |   |
|---|---|
| <input type="checkbox"/> Acetaminophen (Tylenol) 650mg PO x1 dose   | <input type="checkbox"/> Famotidine (Pepcid) 20mg IV x1 dose                |
| <input type="checkbox"/> Diphenhydramine (Benadryl) 25mg PO x1 dose | <input type="checkbox"/> Methylprednisolone (Solu-medrol) 125mg IV x 1 dose |

### Medication Orders

#### Iron Sucrose (Venofer)

- Iron Sucrose 200mg in sodium chloride 0.9% 100mL IV over 30min every 1 to 7 days x \_\_\_ doses
- Iron Sucrose \_\_\_mg in sodium chloride 0.9% \_\_\_mL IV over \_\_\_ min every \_\_\_ days x \_\_\_ doses

#### Ferric Carboxymaltose (Injectafer)

- Ferric Carboxymaltose 750mg in sodium chloride 0.9% 250 mL IV over 30min every 7 days x 2 doses

#### Ferumoxytol (Feraheme)

- Ferumoxytol 510 mg in sodium chloride 0.9% 100mL IV over 30min x 1 dose
- Ferumoxytol 510 mg in sodium chloride 0.9% 100mL IV over 30min every \_\_\_ days x \_\_\_ doses

### MANAGEMENT OF SIDE EFFECTS

If adverse reaction (Hypotensive Reaction of SBP drop 25mmHg, phlebitis/vein spasm, abdominal/leg cramps, nausea, diarrhea): Hold infusion x 30 mins, if symptoms improve resume infusion at half previous rate, if symptoms persist give 500ml IV sodium chloride 0.9% bolus and call provider. If afterhours call hospitalist.

### IF ANAPHYLACTIC REACTION OCCURS

Stop infusion immediately, notify provider and/or Hospitalist on duty if afterhours, administer oxygen prn, administer Benadryl 50 mg IV or IM once STAT, administer Epinephrine (1:1000) 0.5mg IM once STAT, possible admission to emergency department for further evaluation/treatment.

<p><i>Patient Name:</i></p> <p><i>DOB:</i></p>
--



**Ordering Facility/Provider Information**

**By signing below, I affirm the following:**

I am responsible for the care of the patient identified on this form.

I hold an active, unrestricted license to practice medicine in \_\_\_\_\_ (*specify state*)

My physician license number is # \_\_\_\_\_ and I am acting within my scope of practice and authorized by law to order infusion of the medication described above for the patient identified on this form.

<b>Provider Signature:</b> _____	<b>Date</b> _____
<b>Printed name:</b> _____	<b>Phone:</b> _____
<b>Fax:</b> _____	

<b>KVH Provider Co-signature:</b> <i>*Required for all external orders</i>	
<b>Provider Signature:</b> _____	<b>Date</b> _____
<b>Printed name:</b> _____	

<i>Patient Name:</i>
<i>DOB:</i>